



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib



CrossMark

Data Article

Data on copper level in the blood of patients with normal and abnormal angiography

Leila Amiri^a, Ali Movahed^b, Dariush Iranpour^c,
Afshin Ostovar^d, Alireza Raeisi^d, Mozhgan Keshtkar^a,
Najmeh Hajian^b, Sina Dobaradaran^{a,e,f,*}

^a Department of Environmental Health Engineering, Faculty of Health, Bushehr University of Medical Sciences, Bushehr, Iran

^b Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, Biochemistry Group, Bushehr, Iran

^c Bushehr University of Medical Sciences, Department of Cardiology, Faculty of Medicine, Bushehr, Iran

^d The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

^e The Persian Gulf Marine Biotechnology Research Center, The Persian Gulf Biomedical Sciences Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

^f Systems Environmental Health, Oil, Gas and Energy Research Center, The Persian Gulf Biomedical Sciences Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

ARTICLE INFO

Article history:

Received 19 July 2016

Received in revised form

2 August 2016

Accepted 9 August 2016

Available online 18 August 2016

Keywords:

Angiography

Cardiovascular disease

Copper

Bushehr

ABSTRACT

In this data article, we measured the levels of copper in the blood of patients undergoing coronary angiography. The samples were taken from patients with cardiovascular disease in Bushehr's university hospital, Iran. Patients were divided in two groups: normal angiography and abnormal angiography. After the chemical digestion of samples, the concentration levels of Cu in both groups were determined by using inductively coupled plasma optical spectrometry (ICP-OES).

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Correspondence to: The Persian Gulf Marine Biotechnology Research Center, Boostan 19 Alley, Imam Khomeini Street, Bushehr, Iran. Fax: +98 7514763448.

E-mail address: s.dobaradaran@bpums.ac.ir (S. Dobaradaran).

<http://dx.doi.org/10.1016/j.dib.2016.08.021>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Medicine
More specific subject area	Heart, biochemistry heart
Type of data	Table
How data was acquired	ICP-OES (SPECTRO (Germany), Spectro arcos)
Data format	Preliminary data, analyzed
Experimental factors	The samples were from patients with cardiovascular disease, with normal and abnormal angiography. After sampling, the blood serum was isolated by centrifugation. Then they were stored at -80°C until final analysis
Experimental features	Determine the level of copper concentration level in blood samples of patients with cardiovascular disease
Data source location	Bushehr, Iran
Data accessibility	Data is with this article.

Value of the data

- The data can be used in other studies to determine trace elements in the blood.
- The data can be used to evaluate the effect of trace elements on heart function.
- The data can be used for further studies on the trace element effects on other human organs as well as the possible role of trace elements in preventing diseases.

1. Data

We measured concentration levels of Cu in blood samples from patients with cardiovascular disease. General data on patients involved in present study is shown in Table 1. The concentration levels of Cu in patients with normal angiography were in the range of 24.01–135.76 $\mu\text{g}/\text{dL}$ with a mean concentration level of 54.14 $\mu\text{g}/\text{dL}$. The concentration levels of Cu in patients with abnormal angiography were in the range of 33.6–221.3 $\mu\text{g}/\text{dL}$ with a mean concentration level of 60.9 $\mu\text{g}/\text{dL}$ (Table 2). Chi-Square analysis also showed that between all characteristics shown in Table 1, only age had statistically significant relationship with angiography type (p -value=0.033).

2. Experimental design, materials and methods

Blood samples were taken from a total of 120 patients, including the patients with normal angiography ($N=60$) and patients with abnormal angiography ($N=60$). These individuals were randomly selected from cardiovascular patients undergoing coronary angiography, at university heart hospital of Bushehr. ECG, echo and angiography were performed for both groups. These who had normal ECG, echo and normal angiography, were placed in the control group, and so people with ECG, echo and abnormal angiogram were classified in the category of abnormal. After sampling, the blood serum was isolated by centrifuge. Then they were stored at -80°C until final analysis. For digestion of samples, 2.5 ml of serum sample was mixed with 2 ml of HNO_3 and 1 ml of H_2O_2 . The resulting mixture was placed in an oven for 2 h at 50°C . After that, 1 ml of concentrated nitric acid and a few drops of H_2O_2 were added to the solution. Then solution was placed on a hot plate, until clear solution was obtained. The resulting clear solution was cooled down at room temperature and 5 ml HNO_3 0.1 N, was added and a volume of 50 ml from sample was prepared by using deionized water [1]. After chemical digestion and preparation, the samples were read by using ICP-OES [2–5].

Table 1

General characteristics of patients with both normal angiography and abnormal angiography.

General characteristics of patients		Angiography	
		Normal	Abnormal
Gender	Male	50.7%	49.3%
	Female	49.0%	51.0%
Age group	< 40 year	87.5%	12.5%
	41–54 year	56.0%	44.0%
	55–64 year	50.9%	49.1%
	65–74 year	28.6%	71.4%
BMI	> 75 year	20.0%	80.0%
	Normal	45.2%	54.8%
	Overweight	52.5%	47.5%
Mitral regurgitation	Obese	50.0%	50.0%
	Positive	43.7%	56.3%
Family history of heart disease	Negative	63.9%	36.1
	Yes	48.1%	51.9%
Duration of having heart disease	No	53.8%	46.2%
	x < 1 year	51.8%	48.2%
	1–5 year	50.0%	50.0%
Hypertension	5–10 year	0.0%	100.0%
	No	35.9%	64.1%
Diabetes	Yes	57.7%	42.3%
	Yes	38.5%	61.5%
Smoking	No	52.7%	47.3%
	Yes	59.2%	40.8%
	No	44.1%	55.9%

Table 2Mean, maximum and minimum concentration level ($\mu\text{g}/\text{dL}$) of Cu in patient with normal angiography and abnormal angiography.

Cu level in blood samples	Normal angiography	Abnormal angiography
Number	60	60
Minimum	24.01	33.6
Maximum	135.76	221.3
Mean	54.14	60.9
Sig.	0.096	0.096

Acknowledgments

The authors are grateful to Bushehr University of Medical Sciences for their financial support (Grant no 20/71/7151) and the laboratory staff of the Environmental Health Engineering Department for their cooperation. The funder had no role in study design, data collection and analysis, or preparation of the manuscript.

Transparency document. Supplementary material

Transparency data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.08.021>.

References

- [1] R. Nawaz, E. Zahir, S. Siddiqui, A. Usmani, K.F. Shad, The role of trace metals and environmental factors in the onset and progression of schizophrenia in Pakistani population, *World J. Neurosci.* 4 (2014) 450.
- [2] H. Arfaeinia, I. Nabipour, A. Ostovar, Z. Asadgol, E. Abuee, M. Keshtkar, S. Dobaradaran, Assessment of sediment quality based on acid-volatile sulfide and simultaneously extracted metals in heavily industrialized area of Asaluyeh, Persian Gulf: concentrations, spatial distributions, and sediment bioavailability/toxicity, *Environ. Sci. Pollut. Res.* 9 (2016) 1–20.
- [3] D.R. Abadi, S. Dobaradaran, I. Nabipour, X. Lamani, M. Ravanipour, R. Tahmasebi, S. Nazmara, Comparative investigation of heavy metal, trace, and macro element contents in commercially valuable fish species harvested off from the Persian Gulf, *Environ. Sci. Pollut. Res.* 22 (2015) 6670–6678.
- [4] S. Dobaradaran, K. Naddafi, S. Nazmara, H. Ghaedi, Heavy metals (Cd, Cu, Ni and Pb) content in two fish species of Persian Gulf in Bushehr Port, Iran., *Afr. J. Bioethanol.* 9 (2013) 6191–6193.
- [5] V.N. Karbasdehi, S. Dobaradaran, I. Nabipour, H. Arfaeinia, R. Mirahmadi, M. Keshtkar, Data on metal contents (As, Ag, Sr, Sn, Sb, and Mo) in sediments and shells of *Trachycardium lacunosum* in the northern part of the Persian Gulf. Data Brief. (<http://dx.doi.org/10.1016/j.dib.2016.06.065>).